



[Incidence of gestational trophoblastic disease in Stockholm County, 1975-1988.](https://arctichealth.org/en/permalink/ahliterature24508)

<https://arctichealth.org/en/permalink/ahliterature24508>

Author: F. Flam
V. Lundström-Lindstedt
L E Rutqvist

Author Affiliation: Department of Obstetrics and Gynecology, Karolinska Hospital, Stockholm, Sweden.

Source: Eur J Epidemiol. 1992 Mar;8(2):173-7

Date: Mar-1992

Language: English

Publication Type: Article

Keywords: Adolescent
Adult
Choriocarcinoma - epidemiology
Female
Humans
Hydatidiform Mole - epidemiology
Incidence
Middle Aged
Pregnancy
Registries
Sweden - epidemiology
Trophoblastic Neoplasms - epidemiology
Uterine Neoplasms - epidemiology
Vital statistics

Abstract: An epidemiological study with the aim of establishing the incidence of hydatidiform mole, persistent trophoblastic disease and choriocarcinoma in Stockholm County was performed. Based on the regional cancer registry and hospital registers, the incidence for 1975-1988 was calculated for the number of deliveries as well as the total number of pregnancies. Of the molar pregnancies, 6% were treated with chemotherapy because of invasive mole or choriocarcinoma. Non-molar choriocarcinoma occurred in 1/33,717 deliveries. Difficulties in assessing the incidence of gestational trophoblastic disease are discussed.

PubMed ID: 1322822 [View in PubMed](#)